

Viscous Cosmology and Thermodynamics of Apparent Horizon

M. Akbar ^{*} [†]

*Centre for Advanced Mathematics and Physics
National University of Sciences and Technology
Peshawar Road, Rawalpindi, Pakistan*

Abstract

It is shown that the differential form of Friedmann equations of FRW universe can be recast as a similar form of the first law $T_A dS_A = dE + WdV$, of thermodynamics at the apparent horizon of FRW universe filled with the viscous fluid. It is also shown that the generalized second law of thermodynamics holds at the apparent horizon of FRW universe and preserves dominant energy condition.

PACS numbers: 04.70.Dy, 97.60.Lf

^{*}Email address: ak64bar@yahoo.com

[†]Email address: makbar@camp.edu.pk

Introduction: In the cosmological setting, one can associate Hawking temperature and entropy with the apparent horizon analogous to the hawking temperature and entropy associated with the black hole horizon [1, 2, 3]. In the case of de Sitter space, the event horizon matches with the apparent horizon of FRW universe with $k = 0$ however for more general cosmological models, the event horizon may not exist but the apparent horizon associated with the Hawking temperature and entropy always exists. The thermodynamic properties associated with the apparent horizon of FRW universe filled with the perfect fluid has been studied by many authors (see for examples [1, 2, 4, 6, 5, 7, 8]). The extension of this connection between thermodynamics and gravity has also been carried out in the braneworld cosmology [9, 10, 11, 12]. For a general static spherically symmetric and stationary axisymmetric spacetimes, It was shown that the Einstein field equations can be rewritten [18, 19, 20] as a first law of thermodynamics. More recently, by considering a masslike function, it has been shown [13] that the equilibrium thermodynamics should exist in the extended theories of gravity and the Friedman equations in various theories of gravity can be rewritten as a first law $TdS = dE$ at the apparent horizon of FRW universe. More recently, Cai et al [5] has shown that by employing Clausius relation, $\delta Q = TdS$, to the apparent horizon of a FRW universe, they are able to derive the modified Friedmann equation by using quantum corrected entropy-area relation.

The cosmological models with perfect cosmic fluid has been studied widely in literature however viscous cosmic fluid came much later in the study of the universe [17]. It has been revealed by Barrow [14](also see [15]) that the thermodynamic entropy associated with bulk viscosity violates the dominant energy condition and can decrease very well. This fact was resumed [16] in the framework of the solutions given in reference [15]. By taking into account the entropy of the sources only, it has been deduced that the entropy of the bulk viscous sources cannot decrease since it would violate the second law of thermodynamics for large cosmic time values. However, in order to state the generalized second law, one has to include both the entropy connected with the sources and the entropy associated with background geometry. The most of the study of the second law in the cosmological context has been carried out at the event horizon (see for examples [15, 16]) however the validity of second law within the apparent horizon of FRW universe is of great important to investigate. In this letter we will also investigate this issue. The cosmological models with bulk viscosity have been studied in the literature from various point of view (see for examples [15, 21, 22]).

The purpose of this letter is twofold. The first is to show that the Friedmann equations of FRW universe pertaining cosmic bulk viscosity can be rewritten as a similar form of the first law of thermodynamics. The other is to discuss the generalized second law within the apparent horizon of FRW universe.

Let us start with a FRW universe of metric

$$ds^2 = -dt^2 + a^2(t)\left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2\right), \quad (1)$$

where $d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$ stands for the line element of 2-dimensional unit sphere and the spatial curvature constant $k = +1, 0$ and -1 represents a closed, flat and open universe, respectively. The above metric (1) can be rewritten in spherical form

$$ds^2 = h_{ab}dx^a dx^b + \tilde{r}^2 d\Omega^2, \quad (2)$$

where $\tilde{r} = a(t)r$, $x^0 = t$, $x^1 = r$ and $h_{ab} = \text{diag}(-1, \frac{a^2}{1-kr^2})$. One can work out the dynamical apparent horizon from the relation $h^{ab}\partial_a\tilde{r}\partial_b\tilde{r} = 0$ which turns out $\frac{1}{\tilde{r}_A^2} = H^2 + k/a^2$, where $H \equiv \frac{\dot{a}}{a}$ is the Hubble parameter and the dots denote derivatives with respect to the cosmic time. let the universe be filled with a viscous fluid and $u^\mu = (u^0, u^i)$ is the four-velocity of the fluid. In comoving coordinates $u^0 = 1$ and $u^i = 0$. Define $h_{\mu\nu} = g_{\mu\nu} + u_\mu u_\nu$, $w_{\mu\nu} = h_\mu^\alpha h_\nu^\beta u_{[\alpha;\beta]}$ and $\theta_{\mu\nu} = h_\mu^\alpha h_\nu^\beta u_{(\alpha;\beta)}$ as a projection tensor, rotation tensor and expansion tensor, respectively. The scalar expansion is $\theta \equiv \theta^\mu_\mu = u^\mu_{;\mu}$ and $\sigma_{\mu\nu} = \theta_{\mu\nu} - \frac{1}{3}h_{\mu\nu}\theta$ is the shear tensor. Taking into consideration of metric (1), $u^\mu_{;\nu} = h^\mu_\nu \frac{\dot{a}}{a}$ which implies that the rotation and shear tensors vanish, i.e. $w_{\mu\nu} = \sigma_{\mu\nu} = 0$. While the scalar expansion is $\theta = 3\frac{\dot{a}}{a} = 3H$. Hence the energy-momentum tensor of viscous fluid in the background of metric (1) finally becomes [21]

$$T_{\mu\nu} = (\rho + P - \zeta\theta)u_\mu u_\nu + (P - \zeta\theta)g_{\mu\nu}, \quad (3)$$

where ρ , P and ζ are the energy density, thermodynamic pressure and bulk viscosity respectively. The components of energy-momentum tensor are $T_{00} = \rho$, $T_{0i} = 0$ and $T_{ij} = (P - \zeta\theta)g_{ij}$. Therefore, the total effect of the bulk viscosity is to reduce the pressure P of the perfect fluid by an amount $\zeta\theta$, so that the effective pressure of the viscous fluid turns out to be $\tilde{P} = P - \zeta\theta$. The energy conservation $T^\mu_{;\nu} = 0$, yields $\dot{\rho} + 3H(\rho + \tilde{P}) = 0$. Now we turn to discuss the thermodynamic interpretation of Friedman equations at the apparent horizon of FWR universe. For this purpose, we first define the surface gravity $\kappa = \frac{1}{2\sqrt{-h}}\partial_a(\sqrt{-h}h^{ab}\partial_b\tilde{r})$ at the apparent of FRW universe. By utilizing above relation, one can easily find that the surface gravity at the apparent horizon of FRW universe yields

$$\kappa = \frac{-1}{\tilde{r}_A}(1 - \frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A}). \quad (4)$$

When $\frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A} \leq 1$ implies the surface gravity $\kappa \leq 0$ which leads to the temperature $T = \kappa/2\pi \leq 0$. However, in reference [1], an approximation, $\frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A} \ll 1$, has been used while determining horizon temperature. They found $T = \frac{1}{2\pi\tilde{r}_A}$ by approximating surface gravity $|\kappa| = \frac{1}{\tilde{r}_A}$. Now we turn to define energy of the universe within the apparent horizon. We take total matter energy $E = V\rho$ inside a sphere of radius $\tilde{r}$ which is also the Misner-Sharp energy [25], $E = \frac{\tilde{r}}{2}(1 - h^{ab}\partial_a\tilde{r}\partial_b\tilde{r})$, within the apparent horizon. Since at the apparent horizon ($h^{ab}\partial_a\tilde{r}\partial_b\tilde{r} = 0$), so the Misner-Sharp energy is in fact total matter energy inside the sphere of radius $\tilde{r}_A$ and is given by

$$E = V\rho. \quad (5)$$

The entropy $S = A/4$ is the horizon entropy, where $A = 4\pi\tilde{r}_A^2$ is the horizon area. Note that we use the units $\hbar = c = G = \kappa_B = 1$. From the Einstein field equations $G_{\mu\nu} = 8\pi T_{\mu\nu}$, the Friedman equations for the viscous fluid of stress-energy tensor (3) can be written as

$$H^2 + \frac{k}{a^2} = \frac{8\pi}{3}\rho, \quad (6)$$

$$\dot{H} - \frac{k}{a^2} = -4\pi(\rho + \tilde{P}). \quad (7)$$

In terms of the apparent horizon, the Friedman equation (6) can be rewritten as

$$\frac{1}{\tilde{r}_A^2} = \frac{8\pi}{3}\rho. \quad (8)$$

Let the apparent horizon surface acts as the boundary of the thermal system of the FRW universe. In general, the apparent horizon is not constant but changes with time. Let $d\tilde{r}_A$ be an infinitesimal change in radius of the apparent horizon in time interval dt . This change in the apparent horizon will cause a small change dV in the volume V of the universe. This constitutes two spherical system of space-time with radii $\tilde{r}_A$ and $\tilde{r}_A + d\tilde{r}_A$ with a common source of fluid with non-zero pressure and energy density near horizon. Each space-time constituting a thermal system and satisfying Einstein's equations, differs infinitesimally in the extensive variables volume, energy and entropy by dV , dE and dS , respectively, while having the same values of intensive variables temperature T and pressure $\tilde{P}$. Thus for these two thermal states of the space-time, there must exist a certain relation connecting these thermodynamic quantities. To establish connection among the thermal quantities and Friedman equations, we differentiate equation (8) which implies

$$\frac{1}{\tilde{r}_A^3} d\tilde{r}_A = 4\pi(\rho + \tilde{P})Hdt \quad (9)$$

Multiplying both sides of this equation with a factor $\tilde{r}_A^3(1 - \frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A})$, one can rewrite this equation in a form

$$\frac{\kappa}{2\pi} \frac{d(4\pi\tilde{r}_A^2)}{4} = -4\pi\tilde{r}_A^2(\rho + P)H(1 - \frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A})dt \quad (10)$$

One can recognize that the quantities $\frac{\kappa}{2\pi}$ and $\frac{4\pi\tilde{r}_A^2}{4}$ are the temperature T and entropy S respectively. Therefore the above equation can be rewritten as

$$TdS = -4\pi\tilde{r}_A^3(\rho + \tilde{P})H(1 - \frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A})dt \quad (11)$$

Now we turn to the total matter energy (5) and taking differential of it, one gets

$$dE = 4\pi\tilde{r}_A^2\rho d\tilde{r}_A - 4\pi\tilde{r}_A^3(\rho + \tilde{P})Hdt. \quad (12)$$

Using equations (11) and (12) one yields

$$dE = TdS + WdV, \quad (13)$$

where $W = \frac{1}{2}(\rho - \tilde{P})$ is the work density. The above equation is the unified first law [24] of relativistic thermodynamics. Note that the above thermal identity is obtained from the Friedman equation with viscous fluid together with the characteristics of the apparent horizon, while the author of reference [24] studied the thermodynamics of trapping horizon of dynamical black hole. It has been found that the entropy associated with apparent horizon is proportional to the horizon area which is originally initiated from the black hole horizon entropy that satisfies the so-called area formula [26]. In the case of perfect fluid, $\zeta = 0$, one can compare the above thermal identity with the standard form of the first law, $dE = TdS - PdV$, of thermodynamics. In fact, the negative pressure term $-P$ in the first law is taken the place of the work density W . Notice that for pure de Sitter spacetime, $\rho = -P$, then one acquires the standard form $dE = TdS - PdV$. Note that we are considering the universe as a thermal object with apparent horizon as its boundary. So it useful to define other thermodynamic quantities like Enthalpy H and Gibbs free energy G . The Enthalpy is defined by the equation $H = U + PV$, where $U = E$ is the total internal energy which is taken to be the total matter energy of the universe bounded by the apparent horizon. So the Enthalpy of the universe bounded by the apparent horizon can be written in terms of apparent radius $H = \frac{4\pi}{3}(\rho + P)\tilde{r}_A^3$. So the heat capacity of the universe enveloped by the apparent horizon at constant pressure of the perfect is defined as

$$C_P = \left(\frac{\partial H}{\partial T}\right)_P = \left(\frac{\partial H}{\partial \tilde{r}_A} \frac{\partial \tilde{r}_A}{\partial T}\right)_P, \quad (14)$$

which yields $C_P = 4\pi\tilde{r}_A^2(\rho + P)\left(\frac{\partial \tilde{r}_A}{\partial T}\right)_P$. In general the temperature, $T = \kappa/2\pi \equiv \frac{-1}{2\pi\tilde{r}_A}\left(1 - \frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A}\right)$. However, if one considers the approximation, $\frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A} \ll 1$ (also see [1, 9]), the temperature $T = |\kappa|/2\pi \equiv 1/2\pi\tilde{r}_A$ which implies that the heat capacity of the universe is negative provided the dominant energy condition holds. On the other hand, if one consider general temperature $T = \frac{-1}{2\pi\tilde{r}_A}\left(1 - \frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A}\right)$ associated with the apparent horizon which implies the heat capacity of the universe is positive definite provided $\frac{\dot{\tilde{r}}_A}{H\tilde{r}_A} \leq 1$. It is also interesting to note that the total matter energy $E = V\rho$, entropy $S = A/4$ and temperature $T = 1/2\pi\tilde{r}_A$ satisfy the relation $E = TS$.

Generalized Second Law: Recently a lot of attention has been granted to the generalized second law of thermodynamics in the accelerating universe handled by the dark energy [27]. Using a particular model for dark energy, the generalized second law has been studied in reference [28] with the boundaries of universe enveloped by the apparent as well event horizon. It is important to investigate the generalized second law as defined in the region of the universe bounded by the apparent horizon in more general context.

Let us now consider a region of FRW universe bounded by the apparent horizon filled by a perfect fluid of energy density ρ and pressure P . We assume that the region bounded by the apparent horizon acts as a thermal system with boundary defined by the apparent horizon. Since the apparent horizon is not constant but varies with time. As the apparent radius changes, the volume enveloped by the apparent horizon will also change, however the thermal system

bounded by the apparent horizon remains in equilibrium when it moves from one state to another so that the temperature of the system must be uniform and the same as the temperature of its surroundings. This requires that the temperature of total energy inside the apparent horizon should be in equilibrium with the temperature associated with the apparent horizon because we are not considering the flow of energy through the horizon.

In order to state that a given cosmological solution satisfies/violates the second law of thermodynamics one should define first what is the generalization of the second law to the case of FRW universe. It should include both the entropy due to sources and the entropy associated with the background geometry. So the generalized second law can be expressed as

$$\dot{S}_A + \dot{S}_m \geq 0, \quad (15)$$

where $S_A = A/4G$ is the Bekenstein-Hawking entropy associated with the apparent horizon of FRW universe and S_m is the entropy due to the matter sources inside the apparent horizon. From the Friedmann equations (6) and (7), it can easily be shown that $\dot{\tilde{r}}_A = 4\pi\tilde{r}_A^3 H(\rho + P) \geq 0$ provided the universe driven by the source of perfect fluid preserves dominant energy condition. Since we are assuming the region of a FRW universe enveloped by the apparent horizon as thermal system so that the change of energy dE of the universe from one state to another must be connected through the Friedmann equations together with the characteristics of the apparent horizon. It has been shown [9, 3, 10] that the Friedmann equations at apparent horizon of FRW universe filled with perfect fluid satisfied the thermal identity $dE = T_A dS_A + W dV$ instead of satisfying the standard first law $T_A dS_A = dE + P dV$, where T_A is the temperature associated with the apparent horizon. Since we are assuming the local equilibrium so that the change of matter energy dE_m satisfies the first law of thermodynamics, $dE_m = T_A dS_m - P dV$, where $T_A = |\kappa|/2\pi$ is the temperature associated with the apparent horizon and is given by

$$T = \frac{1}{2\pi\tilde{r}_A} \left(1 - \frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A}\right), \quad (16)$$

where $\frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A} \leq 1$ ensures that the temperature is positive. From the first law, $dE_m = T_A dS_m - P dV$, one can obtain

$$T_A \dot{S}_m = 4\pi\tilde{r}_A^2 \dot{\tilde{r}}_A (\rho + P) - 4\pi\tilde{r}_A^3 H(\rho + P). \quad (17)$$

Now we turn to find out the entropy change $\dot{S}_H$ associated with the apparent horizon of FRW universe which yields

$$T_A \dot{S}_A = 4\pi\tilde{r}_A^3 H(\rho + P) - 2\pi\tilde{r}_A^2 (\rho + P) \dot{\tilde{r}}_A. \quad (18)$$

Adding equations (17) and (18), one gets the expression for the generalized second law

$$T_A (\dot{S}_m + \dot{S}_H) = 2\pi\tilde{r}_A^2 \dot{\tilde{r}}_A (\rho + P). \quad (19)$$

It is obvious from the above equation that $\dot{S}_m + \dot{S}_H \geq 0$ implies that the generalized second law holds provided $\rho + P \geq 0$.

Conclusion: It is shown that the differential form of Friedmann equations of FRW universe filled with a viscous fluid can be rewritten as a similar form of the first law $TdS = dE + WdV$, of thermodynamics at the apparent horizon of FRW universe. It can easily be seen that the similar identity also holds for the perfect fluid when bulk viscosity of the fluid is zero. It is also shown that the heat capacity of the universe filled with perfect fluid is negative if one considers the approximate temperature $T = |\kappa|/2\pi = 1/2\pi\tilde{r}_A$. However if one utilizes the general expression $T = \frac{-1}{2\pi\tilde{r}_A}(1 - \frac{\dot{\tilde{r}}_A}{2H\tilde{r}_A})$, the heat capacity of the universe is positive definite provided $\frac{\dot{\tilde{r}}_A}{H\tilde{r}_A} \leq 1$ which implies that the universe within the apparent horizon of FRW universe is thermodynamically stable. We also verify that the generalized second law of thermodynamics at the apparent horizon of FRW universe holds provided that the matter source satisfies the dominant energy condition. It is interesting to investigate the generalized second law for the extended theories of gravity in a region bounded by the apparent horizon of FRW universe. The work in this respect is under progress.

Acknowledgments

I would like to thank Rong-Gen Cai for his useful comments. The work is supported by a grant from National university of Sciences and Technology.

References

- [1] R. G. Cai and S. P. Kim, JHEP **0502**, 050 (2005)[arXiv:hep-th/0501055]
- [2] G. W. Gibbons and S. W. Hawking, Phys.Rev. D**15**, 2738 (1977).
- [3] M. Akbar and R. G. Cai, Phys. Rev. D**75**, 081003(2007)
- [4] C. Eling, R. Guedens and T. Jacobson, Phys. Rev. Lett. **96**,121301(2006).
- [5] R. G. Cai, L. M. Cao, and Y. P. Hu, hep-th/0807.1232
- [6] M. Akbar and R. G. Cai, Phys. Lett. B**648**, 243(2007)
- [7] R. G. Cai and L. M. Cao, Phys. Rev. D**75**, 064008(2007)
- [8] M. Akbar and R. G. Cai, Phys. Lett. B**635**, 7 (2006). [arXiv:hep-th/0602156]
- [9] R. G. Cai and L. M. Cao, hep-th/0612144
- [10] A. Sheykhi, B. Wang and R. G. Cai, Nucl. Phys. B**779**, 1(2007) [hep-th/0701198]
- [11] A. Sheykhi, B. Wang and R. G. Cai, hep-th/0701261

- [12] R. G. Cai, hep-th/0712.2142
- [13] Y. Gong and A. Wang, hep-th/0704.0793
- [14] J. D. Barrow, Nucl. Phys. **B310**, 743(1988)
- [15] M. Giovannini, Phys. Rev. D **59**, 121301(1999)
- [16] W. Zimdahl and D. Pavon, Phys. Rev. D **61**, 108301(2000)
- [17] C. W. Misner, Astrophys. J. **151**, 431(1968)
- [18] D. kothawala, S. Sarkar, and T. Padmanabhan, arXiv:gr-qc/0701002.
- [19] T. Padmanabhan, Class. Quant. Grav. **19**, 5387 (2002)[arXiv: gr-qc/0204019]; Phys. Rept. **406**, 49 (2005)[arXiv: gr-qc/0311036].
- [20] A. Paranjape, S. Sarkar and T. Padmanabhan, Phys. Rev. D **74**, 104015(2006)[arXiv:hep-th/0607240].
- [21] Iver Brevik, gr-qc/0404095
- [22] W. Zimdahl, Phys. Rev. D **53**, 5483(1996)
- [23] M. Akbar and R. G. Cai, Phys. Rev. D **75**, 084003(2007); M. Akbar and R. G. Cai, Phys. Lett. **B635**, 7(2006)[hep-th/0602156]; M. Akbar and R. G. Cai, Phys. Lett. **B648**, 243(2007)[gr-qc/0612089]; A. V. Frolov and L. Kofman, JCAP **0305**, 009 (2003); R. Bousso, Phys. Rev. D **71**, 064024(2005)[arXiv: hep-th/0412197];
- [24] S. A. Hayward, S. Mukohyana, and M. C. Ashworth, Phys. Lett. A **256**, 347(1999); S. A. Hayward, Class. Quant. Grav. **15**, 3147(1998)
- [25] C. W. Misner and D. H. Sharp, Phys. Rev. **136**, B571(1964)
- [26] R. M. Wald, Phys. Rev. D **48**, 3427(1993)
- [27] G. Izquierdo and D. Pavon, Phys. Lett. **B633**, 420(2006); G. Izquierdo and D. Pavon, Phys. Lett. **B639**, 1(2006); M. R. Setare, JCAP **0701**, 023(2007); H. Mohseni Sadjadi, Phys. Lett. **B645**, 108(2007); Y. Gong, B. Wang and A. Wang, JCAP **0701**, 024(2007); J. Zhou, B. Wang, Y. Gong and E. Abdalla, gr-qc/0705.1264
- [28] B. Wang, Y. Gong and E. Abdalla, Phys. Rev. D **74**, 083520(2006)